

Nucleon axial charge from quenched lattice QCD with domain wall fermions and DBW2 gauge action

Shigemi Ohta^{ab} for the RBC collaboration*

^aInstitute of Particle and Nuclear Studies, KEK, Tsukuba, Ibaraki 305-0801, Japan

^bRIKEN-BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA

The domain wall fermion (DWF) method, with its almost perfectly preserved chiral symmetry on the lattice, makes the calculation of the nucleon axial charge particularly easy. By maintaining chiral symmetry and using the Ward-Takahashi (WT) identity, one has $Z_A = Z_V$ and the bare lattice calculation yields the physical value without explicit renormalization. The DBW2 improved gauge action provides further enhancement of the symmetry and hence a more accurate WT identity at coarse lattice spacing. Taking advantage of these methods, we confirmed a significant volume dependence of the nucleon axial charge on $(1.2\text{fm})^3$ and $(2.4\text{fm})^3$ lattice volumes.

Domain wall fermions (DWF) [1] preserve chiral symmetry on the lattice by introducing a fictitious fifth dimension in which the symmetry violation is exponentially suppressed. The DBW2 (“doubly blocked Wilson 2”) improved gauge action [2] improves the approach to the continuum by adding rectangular (2×1) Wilson loops to the action. By combining the two, the “residual mass,” which controls low energy chiral behavior, is driven to as small as $am_{\text{res}} < O(10^{-4})$, or $\ll 1\text{MeV}$. Successful numerical lattice calculations investigating chiral symmetry and the ground-state hadron mass spectrum [3], Kaon matrix elements [4], and negative-parity excited nucleon (N^*) mass [5] have been reported. Here we apply the method to the axial charge, g_A , the simplest of nucleon electroweak matrix elements.

From neutron β decay, we know $g_V = G_F \cos \theta_c$ and $g_A/g_V = 1.2670(35)$ [6]. These quantities are defined as $q^2 \rightarrow 0$ limits of the form factors $g_V(q^2)$ and $g_A(q^2)$ in the relevant vector and axial current matrix elements, $\langle n | V_\mu^-(x) | p \rangle$,

$$i\bar{u}_n[\gamma_\mu g_V(q^2) + q_\lambda \sigma_{\lambda\mu} g_M(q^2)]u_p e^{-iqx},$$

$$\text{and } \langle n | A_\mu^-(x) | p \rangle,$$

$$i\bar{u}_n \gamma_5 [\gamma_\mu g_A(q^2) + q_\mu g_P(q^2)]u_p e^{-iqx}.$$

On the lattice, in general, we calculate the relevant matrix elements of these currents with a lattice cutoff, $a^{-1} \sim 1\text{--}2\text{ GeV}$, and extrapolate to the continuum, $a \rightarrow 0$, introducing lattice renormalization: $g_{V,A}^{\text{ren}} = Z_{V,A}^{\text{lat}} g_{V,A}^{\text{lat}}$. Also, unwanted lattice artefacts may result in unphysical mixing of chirally distinct operators. DWF make the calculation particularly easy, because the chiral symmetry is almost exact; the relation $Z_A = Z_V$ is maintained, so the lattice ratio $(g_A/g_V)^{\text{lat}}$ directly yields the renormalized ratio.

It is interesting to note the pre-QCD model estimations for this quantity. The non-relativistic quark model in its simplest form gives $5/3$, and the MIT bag model gives 1.07 . The lattice QCD calculations with Wilson or clover fermions underestimate g_A by 10 to 25 % [7]. These calculation also suffer from systematic errors arising from $Z_A \neq Z_V$ and other renormalization complications.

Our formulation follows the standard one, and was summarized in our Lattice 2001 proceedings [8]. In addition to the wall sources, sequential sources are added [9].

Numerical calculations with Wilson (single plaquette) gauge action were performed at $\beta = 6.0$ on a $16^3 \times 32 \times 16$ lattice and $M_5 = 1.8$ with

*Current members are Y. Aoki, T. Blum, N. Christ, M. Creutz, C. Dawson, T. Izubuchi, L. Levkova, X. Liao, G. Liu, R. Mawhinney, Y. Nemoto, J. Noaki, S. Ohta, K. Orginos, S. Prelovsek, S. Sasaki and A. Soni. We thank RIKEN, BNL and the U.S. DOE for providing the facilities essential for the completion of this work.

nucleon source at $t = 5$, sink at 21, and current insertions in between. Results from 400 gauge configurations are summarized as follows. The local vector current renormalization $Z_V = 1/g_V^{\text{lattice}}$ is well-behaved: the value 0.764(2) at $m_f = 0.02$ agrees well with $Z_A = 0.7555(3)$ of the corresponding local axial current renormalization [3,10]. A linear fit gives $Z_V = 0.760(7)$ at $m_f = 0$, and a quadratic fit, 0.761(5). The axial to vector charge ratio, g_A/g_V , averaged in $10 \leq t \leq 16$ time slices, yields a value of 0.81(11) in the linearly extrapolated chiral limit $m_f = 0$. These are essentially the same as last year [8], but have been confirmed with much higher accuracy obtained through higher statistics.

Possible reasons for this small value of g_A/g_V are finite lattice volume [11], excited states (small separation between t_{source} and t_{sink}), and quenching (zero modes, absent pion cloud, and so on). Here we concentrate on the first possibility. To investigate size-dependence, we need good chiral behavior or sufficient proximity to the continuum, and big enough lattice spatial volume at the same time. Improved gauge actions help both [12]. In particular the DBW2 action [2] gives very small residual chiral symmetry breaking, $am_{\text{res}} < 10^{-3}$, at a lattice spacing of $a \sim 0.15$ fm. At this lattice cutoff the hadron mass spectrum is well-behaved in the chiral limit too: $am_\rho = 0.592(9)$, $m_\rho/m_N \sim 0.8$, and $m_\pi(m_f = 0.02) \sim 0.3a^{-1}$.

DBW2 calculations are performed with both wall and sequential sources on two lattices: $8^3 \times 24 \times 16$ ($\sim (1.2\text{fm})^3$), 405 configurations (wall and sequential), and $16^3 \times 32 \times 16$ ($\sim (2.4\text{fm})^3$), 100 configurations (wall) and 204 (sequential). Nucleon source-sink separation is set at about 1.5 fm, and quark mass values of $m_f = 0.02, 0.04, 0.06, 0.08$ and 0.10 are used. So the pion mass is as low as $m_\pi \geq 390\text{MeV}$ and satisfies $m_\pi L \geq 4.8$ and 2.4 , respectively. The residual mass at this cutoff is about 0.8 MeV or $am_{\text{res}} \sim 6 \times 10^{-4}$.

Again the vector and axial current renormalization factors are well-behaved (see Figure 1.) The vector renormalization, Z_V , calculated as $1/g_V$, shows slight quadratic dependence on m_f as expected: $V_\mu^{\text{conserved}} = Z_V V_\mu^{\text{local}} + \mathcal{O}(m_f^2 a^2)$, yielding a value $Z_V = 0.791(17)$ in the chiral limit. This agrees well with the axial renor-




Figure 1. Local current renormalization factors, vector (Z_V , circle) and axial (Z_A , square).

malization, $Z_A = 0.77759(45)$, obtained from $\langle A_\mu^{\text{conserved}}(t)[\bar{q}\gamma_5 q](0) \rangle = Z_A \langle A_\mu^{\text{local}}(t)[\bar{q}\gamma_5 q](0) \rangle$ [10]. This assures that the lattice calculation of the axial-to-vector ratio directly yields its continuum value without any further renormalization: $(g_A/g_V)^{\text{lat}} = (g_A/g_V)^{\text{ren}}$.

In Figure 2 the quark-mass dependence of the ratio obtained from sequential source calculations is summarized for the two lattice volumes: Clear volume dependence is seen between $(2.4\text{fm})^3$ and $(1.2\text{fm})^3$ volumes. The large volume results with the sequential source show a very mild m_f dependence and extrapolate to a value of $(g_A/g_V)^{\text{ren}} = 1.21(3)(+4)$. The first error is statistical and the second is systematic from a preliminary finite-size analysis. At the lightest quark mass value of $m_f=0.02$, the volume dependence is no longer statistically significant. Indeed at this m_f even the larger volume may not be big enough. In Figure 3 the same data are presented in physical units together with the small volume Wilson gauge action results. The smaller volume results with Wilson ($16^3 \times 32$ at $a \sim 0.1$ fm) and DBW2 ($8^3 \times 24$ at $a \sim 0.15$ fm) actions are mutually




Figure 2. $(g_A/g_V)^{\text{ren}}$: m_f and volume dependence. Sequential source results on a large (cross) and small (circle) lattice. Wall source results are consistent with larger errors.

consistent. In summary we confirmed a lattice-volume dependence of the nucleon axial charge, g_A/g_V . Similar dependence was seen in the bare tensor charge. Further investigation is under way.

REFERENCES

1. D.B. Kaplan, Phys. Lett. B288, 342 (1992); Y. Shamir, Nucl. Phys. B406, 90 (1993); V. Furman and Y. Shamir, Nucl. Phys. B439, 54 (1995).
2. P. de Forcrand *et al.* (QCD-TARO collaboration), Nucl. Phys. B577, 263 (2000).
3. T. Blum, *et al.* (RBC collaboration), to appear in Phys. Rev. D, hep-lat/0007038.
4. T. Blum, *et al.* (RBC collaboration), RBRC Scientific Articles 4; hep-lat/0110075, submitted for publication in Phys. Rev. D.
5. S. Sasaki, T. Blum and S. Ohta (for RBC collaboration), Phys. Rev. D65, 074503 (2002).
6. The Particle Data Group.




Figure 3. $(g_A/g_V)^{\text{ren}}$: m_f and volume dependence in physical scale set by ρ meson mass. Wilson (diamond) and DBW2 results are consistent in the smaller volumes.

7. M. Fukugita, Y. Kuramashi, M. Okawa and A. Ukawa, Phys. Rev. Lett. 75, 2092 (1995); K.F. Liu, S.J. Dong, T. Draper and J.M. Wu, Phys. Rev. D49, 4755 (1994); M. Göckeler *et al.*, Phys. Rev. D53, 2317 (1996); D. Dolgov *et al.*, hep-lat/0201021; S. Capitani *et al.*, Nucl. Phys. B (Proc. Suppl.) 79, 548 (1999); R. Horsley *et al.*, Nucl. Phys. B (Proc. Suppl.) 94, 307 (2001); S. Güsken *et al.*, Phys. Rev. D59, 114502 (1999).
8. S. Sasaki, Nucl. Phys. Proc. Suppl. 106, 30 (2002); and references cited there in.
9. K. Orginos for the RBC collaboration, in these proceedings.
10. T. Blum *et al.* (RBC Collaboration), Phys. Rev. D66, 014504 (2002).
11. R.L. Jaffe, Phys. Lett. B529, 105 (2002); T.D. Cohen, Phys. Lett. B529, 50 (2002).
12. RBC collaboration, in preparation.